



1. Owner Information: This information is used not only for the animal record, but is also used for programming microchips. Accurate information is needed to ensure ownership is established correctly.

First Name:	Last Name:	Preferred Pronouns	
Income Verification Statement: I have reviewed the income qualification standards and certify that my household income is less than 80% of the median family income.		Street Address:	City: Zip Code:
<input type="radio"/> Yes <input checked="" type="radio"/> No			
Mobile Phone:	Alternate Phone:	Email:	

2. Animal Information

Name:	Primary Breed:	Size:	
Gender:	Date of Birth (best estimate):	Primary Color:	Secondary Color:
Color Pattern:	Any medical concerns?		Any handling instructions or behavior concerns?

3. Additional Services. Please select any services that you would like to add.

Microchip: \$10

yes no

Distemper (DHPP) Vaccine: \$10

yes no

Rabies Vaccine: \$15

yes no

Bordatella Vaccine: \$10

yes no

E-collar (cone): \$5

yes no

Dewormer (Nemex): \$10

yes no

Flea Treatment (topical, 1 month): \$15

yes no

Ear Cleaning: \$5

yes no

Umbilical Hernia Repair: \$25

yes no

Nail Trim: free

yes no

Are you using a voucher for surgery or any services?

yes no

If yes, please enter the Voucher Number

4. Seattle Humane

Post-Operative Surgery Instructions

[Post-Op Instructional Video](#)

Spay/Neuter Operation

- The neuter operation for males involves surgical removal of both testicles.
- The spay operation for females involves surgical removal of both ovaries and the uterus.

Post-Op Instructions

Food & Water

- Wait at least 2 hours after returning home to give your pet food or water. Remember, anesthesia can make pets nauseous. Animals should be offered $\frac{1}{2}$ their normal meal the evening of surgery. Resume normal feeding the following day and it may take a few days for your pet's appetite to return to normal.
- It is not uncommon for your pet to have a bout of loose stool or to vomit from the stress of surgery, but if it persists please call and let us know or contact your local veterinarian or emergency clinic.

Activity

- Should be confined to a clean, small, warm, and dry area indoors.
- Limit your dog's activity level for the next 10-14 days (i.e. no running, jumping, etc.).
- Leash walks only and try to hinder anything, like jumping or running, that will stretch the incision site.
- Do not allow your dog to be unattended around stairs the night after surgery or the following day. He or she may still be groggy and a little uncoordinated.

Sutures:

- Unless otherwise stated, all dogs have internal sutures that will dissolve. Check the incision twice a day for the next 10-20 days to make certain it is healing and not oozing or swelling. Some bruising is normal. The scrotum may still "look full" for a few weeks but should not be hard, red or painful.

Baths

Do not bathe your pet for the next 10-14 days. Also, do not allow your pet to swim or take long outings in the rain. Keep your pet indoors and dry.

Pain Medication

- Your dog will be sent home with oral pain medication (Carprofen). They will get enough for 5 days and they should start the recommended dosage the morning following surgery. Please make sure you are following the instructions on the bottle to administer their medication appropriately. **Dogs are known to shuffle their food around to avoid any medication in their bowl. It is recommended that you wrap pills in cheese, peanut butter, or stuff them in a hot dog.**
- **Puppies/Small Dogs:** Will be sent home with a liquid pain medication for 5 days.

****If you think your pet seems painful in the days after surgery (whining, hiding, aggression, decreased appetite), please call us for assistance.***

E-Collars

- Animals may lick their incision site to soothe the discomfort. It is recommended that you purchase an e-collar/cone and keep it on them for 10-14 days. It is very important that the e-collar is left on the dog while they are unattended (when you are away from home and overnight).

*****If you notice your pet licking their incision and you don't have an e-collar, please purchase one and place it on your pet ASAP. Damage to incision can lead to serious consequences including infection and incisional failure. They can be purchased from us or at most pet stores or veterinary clinics.***

Tattoos

Your pet will be surgically marked with a green tattoo near the incision as visual proof that your pet has been spayed or neutered. For male cats, this is a small surgical mark on the abdomen.

If you have any questions or concerns, please call the following number or email:

(425) 649-7560 or vets@seattlehumane.org